

Lecturer: Olaf Lechtenfeld
 Assistant: Susha Parameswaran

Problem 1: Spinor Representation of the Lorentz Algebra

The Clifford algebra is given by the anticommutation relations:

$$\{\gamma^\mu, \gamma^\nu\} \equiv \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2\eta^{\mu\nu} \mathbb{1}. \quad (1)$$

The Weyl representation of this algebra is given by the following choice of gamma matrices:

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix}, \quad \text{where } \sigma^\mu = (\mathbb{1}, \sigma^i) \text{ and } \bar{\sigma}^\mu = (\mathbb{1}, -\sigma^i), \quad (2)$$

with σ^i being the Pauli matrices, satisfying $\sigma^i \sigma^j = \delta^{ij} + i\epsilon^{ijk} \sigma^k$ and $(\sigma^i)^\dagger = \sigma^i$.

Dirac realized that given any representation of the Clifford algebra, one can immediately write down a spinor representation of the Lorentz algebra as:

$$\sigma^{\mu\nu} = \frac{i}{4} [\gamma^\mu, \gamma^\nu], \quad (3)$$

which is a spinor representation. The spinor representation of the Lorentz group is then obtained via exponentiation

$$S(\Lambda) = e^{-\frac{i}{2} \omega^{\mu\nu} \sigma_{\mu\nu}}, \quad (4)$$

where $\omega^{\mu\nu}$ is the parameter for infinitesimal Lorentz transformations.

A four-component field ψ that transforms accordingly as

$$\psi(x) \rightarrow S(\Lambda)\psi(\Lambda^{-1}x) \quad (5)$$

under Lorentz transformations is called a Dirac spinor.

- (a) Compute the Lorentz generators $\sigma^{\mu\nu}$ using the Weyl representation of the gamma matrices.
- (b) Is this representation irreducible? What does this imply for the Dirac spinor?
- (c) Is this representation Hermitian? Is the corresponding representation of the Lorentz group $S(\Lambda)$ unitary?
- (d) Write down the explicit matrix representing a rotation through an angle θ about the z -axis, that is taking $\omega^{12} = \theta$ and all other components zero. What is the result for $\theta = 2\pi$?

Problem 2: Weyl spinors, helicity and parity

We can define a “fifth” gamma matrix

$$\gamma^5 = \frac{i}{4!} \epsilon^{\mu\nu\rho\sigma} \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 \gamma^4. \quad (6)$$

where $\epsilon^{\mu\nu\rho\sigma}$ is a totally antisymmetric tensor with $\epsilon^{0123} = 1$. It is easy to show that $\{\gamma^5, \gamma^\mu\} = 0$.

Left- and right-handed spinors are defined by the following conditions

$$\gamma^5 \psi_L = -\psi_L, \quad \gamma^5 \psi_R = \psi_R. \quad (7)$$

- (a) Show that an arbitrary Dirac spinor can be split into a left-handed and right-handed part as:

$$\psi = \psi_L + \psi_R, \quad \psi_{L,R} = P_{L,R} \psi, \quad P_{L,R} = \frac{1}{2}(\mathbb{1} \mp \gamma^5) \quad (8)$$

Hint: first show that $(\gamma^5)^2 = \mathbb{1}$.

- (b) Compute γ^5 , $P_{L,R}$ and ψ in the Weyl representation, and show how $\psi_{L,R}$ transform under Lorentz transformations.
- (c) Write down the Dirac Lagrangian in terms of the Weyl spinors $\psi_{L,R}$.
- (d) Show that in the massless case $\psi_{L,R}$ are eigenstates of the helicity operator

$$h = \frac{\vec{p} \cdot \vec{\sigma}}{|\vec{p}|} \quad (9)$$

What are the eigenvalues?

Hint: use the Dirac equation in the Weyl representation.

- (e) In the Standard Model of Particle Physics, the fields mediating the weak interaction, the W^\pm , Z bosons, couple only to the left-handed components of spin 1/2 fields. E.g. the vector Z boson, has a coupling of the form

$$Z^\mu \bar{\psi}_L \gamma_\mu \psi_L = Z^\mu \bar{\psi} \gamma_\mu \frac{1}{2}(\mathbb{1} - \gamma^5) \psi \quad (10)$$

in the Lagrangian. Does this interaction respect parity, $(t, \vec{x}) \rightarrow (t, -\vec{x})$?

Hint: Show that $\psi(t, \vec{x}) = \gamma^0 \psi(t, -\vec{x})$, i.e. parity interchanges left- and right-handed spinors.