

Quantum Field Theory: Exercise Session 1

Lecturer: Olaf Lechtenfeld

Problem 1: Poincaré algebra for the real scalar field

The real Klein-Gordon field $\phi(x)$ is governed by the Lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2} \phi^2. \quad (1)$$

We can write $\phi(x)$ and its conjugate momentum $\pi(x)$ in terms of the creation and annihilation operators $a_{\vec{k}}$ and $a_{\vec{k}}^\dagger$ as

$$\phi(\vec{x}) = \int \tilde{d}k \left[a_{\vec{k}} e^{i\vec{k}\cdot\vec{x}} + a_{\vec{k}}^\dagger e^{-i\vec{k}\cdot\vec{x}} \right], \quad (2)$$

$$\pi(\vec{x}) = -i \int \tilde{d}k \omega_k \left[a_{\vec{k}} e^{i\vec{k}\cdot\vec{x}} - a_{\vec{k}}^\dagger e^{-i\vec{k}\cdot\vec{x}} \right], \quad (3)$$

where

$$\tilde{d}k \equiv \frac{d^3k}{(2\pi)^3 2\omega_k}, \quad \omega_k \equiv \sqrt{\vec{k}^2 + m^2}, \quad \vec{k} \cdot \vec{x} \equiv k^i x^j \delta_{ij}. \quad (4)$$

In the lecture, you have seen that energy and momentum may be written in terms of $a_{\vec{k}}$ and $a_{\vec{k}}^\dagger$ as

$$\begin{aligned} P^0 \equiv H &\equiv \int d^3x T^{00} = \int d^3x \frac{1}{2} \left(\pi^2 + (\vec{\nabla}\phi)^2 + m^2\phi^2 \right) \\ &= \int \tilde{d}k \omega_k a_{\vec{k}}^\dagger a_{\vec{k}}, \end{aligned} \quad (5)$$

and

$$P^i \equiv \int d^3x T^{0i} = \int d^3x \pi(x) \nabla^i \phi(x) = \int \tilde{d}k k^i a_{\vec{k}}^\dagger a_{\vec{k}}, \quad (6)$$

where $T^{\mu\nu}$ is the energy-momentum tensor. The Lorentz generators are given by

$$M^{\mu\nu} \equiv \int d^3x (x^\mu T^{0\nu} - x^\nu T^{0\mu}). \quad (7)$$

- (a) Use the above expressions to write down the boost and rotation generators M_{i0} and M_{ij} in terms of $\phi(x)$ and $\pi(x)$.

- (b) Use the Fourier expansions of $\phi(x)$ and $\pi(x)$ to express the rotation generators M_{ij} in terms of $a_{\vec{k}}$ and $a_{\vec{k}}^\dagger$.

Hint:

$$\int d^3x x_i e^{i\vec{k}\cdot\vec{x}} = \int d^3x \left(-i \frac{\partial}{\partial k_i} \right) e^{i\vec{k}\cdot\vec{x}} = -i(2\pi)^3 \frac{\partial}{\partial k_i} \delta^3(\vec{k}). \quad (8)$$

- (c) Compute the commutators $[P^i, \phi(x)]$ and $[M^{ij}, \phi(x)]$ in terms of $\phi(x)$, with the help of the commutator relations for $a_{\vec{k}}$ and $a_{\vec{k}}^\dagger$.
- (d) *Optional.* Repeat exercises (b) and (c) for the boost generators M_{0i} to find

$$M_{i0} = -i \int \tilde{d}k \omega_k a_{\vec{k}}^\dagger \partial_{k_i} a_{\vec{k}} \quad (9)$$

and compute the commutator $[M^{i0}, \phi(x)]$. Check that the commutator of M_{i0} with M_{j0} satisfies the Lorentz algebra

$$[M^{\sigma\tau}, M^{\alpha\beta}] = i(\eta^{\tau\alpha} M^{\sigma\beta} + \eta^{\sigma\beta} M^{\tau\alpha} - \eta^{\sigma\alpha} M^{\tau\beta} - \eta^{\tau\beta} M^{\sigma\alpha}). \quad (10)$$

Problem 2: The complex scalar field

The free complex Klein-Gordon scalar field is governed by the Lagrangian density

$$\mathcal{L} = \partial_\mu \phi^\dagger \partial^\mu \phi - m^2 \phi^\dagger \phi. \quad (11)$$

Since the complex scalar field carries two degrees of freedom, quantizing it gives rise to two independent creation operators. The mode expansion for ϕ is

$$\phi(\vec{x}) = \int \tilde{d}k \left[a_{\vec{k}} e^{i\vec{k}\cdot\vec{x}} + b_{\vec{k}}^\dagger e^{-i\vec{k}\cdot\vec{x}} \right], \quad (12)$$

where the operators $a_{\vec{k}}$ and $b_{\vec{k}}$ satisfy the commutation relations:

$$[a_{\vec{k}}, a_{\vec{p}}^\dagger] = [b_{\vec{k}}, b_{\vec{p}}^\dagger] = \tilde{\delta}(\vec{k} - \vec{p}) \quad (13)$$

with all other commutators vanishing. The creation operators $a_{\vec{k}}^\dagger$ and $b_{\vec{k}}^\dagger$ create two types of particle, both of mass m and spin zero, which are interpreted as particles and anti-particles.

Notice that \mathcal{L} is invariant under the rigid phase transformation $\phi \rightarrow e^{i\alpha} \phi$. Associated to this symmetry, Noether's theorem gives the conserved charge

$$Q = i \int d^3x (\partial_0 \phi^\dagger \phi - \phi^\dagger \partial_0 \phi). \quad (14)$$

- (a) Write down the mode expansion for ϕ^\dagger and the conjugate momenta, π, π^\dagger .
- (b) Express H and Q in terms of the creation and annihilation operators. Show that $[H, Q] = 0$ and give the interpretation of Q . Comment also on the implications that the theory is free.