

Nonmeromorphic operator product expansion and C_2 -cofiniteness for a family of \mathcal{W} -algebras

Nils Carqueville* and Michael Flohr†

*Physikalisches Institut
University of Bonn
Nussallee 12
53115 Bonn, Germany*

Abstract

We prove the existence and associativity of the nonmeromorphic operator product expansion for an infinite family of vertex operator algebras, the triplet \mathcal{W} -algebras, using results from $P(z)$ -tensor product theory. While doing this, we also show that all these vertex operator algebras are C_2 -cofinite.

1 Introduction

The notion of operator product expansion is fundamental in (quantum) field theory; this notion was originally introduced by Wilson [W] and Kadanoff [Kad]. Physically, it describes the short distance behaviour of the product of two quantum fields $\Phi_1(z_1)$ and $\Phi_2(z_2)$ when it is evaluated near the singularity at $z_1 = z_2$ and therefore encodes part of the local structure of the theory. The singularity arises because the fields are distributions.

In particular, products of fields occur in correlation functions, which eventually allow to compute observables which can then be compared with experimental data. One important advantage of operator product expansion, viewed as a tool that expands the product of *two* fields into a series in which each summand involves only *one* field, is that in this way n -point functions can be expressed in terms of $(n - 1)$ -point functions. This does not only tremendously facilitate concrete computations, but it also structures the theory conceptually.

*nils@th.physik.uni-bonn.de

†flohr@th.physik.uni-bonn.de

While for generic spacetime dimensions a rigorous formulation and treatment of operator product expansion is very difficult, in two-dimensional conformal field theory the situation is much clearer (which is one of many reasons to study conformal field theory, apart from its prominent role and successful application in string theory and statistical physics).

A natural framework to deal with two-dimensional conformal field theory is that of vertex operator algebras [FHL], [LL] and related concepts. In this language, the operator product expansion of fields operating on a *fixed* module (or representation) W for a vertex operator algebra V is expressed in the well-known result

$$\iota_{12}^{-1} \langle w', Y(u, x_1)Y(v, x_2)w \rangle = \left(\iota_{20}^{-1} \langle w', Y(Y(u, x_0)v, x_2)w \rangle \right) \Big|_{x_0=x_1-x_2}, \quad (1.1)$$

where $u, v \in V$, $w \in W$, $w' \in W' = \coprod_{n \in \mathbb{Z}} (W_{[n]})^*$, $\langle \cdot, \cdot \rangle$ denotes the natural pairing between W' and W , and ι_{ij} denotes the operation of expanding a function of x_i and x_j such that only finitely many negative powers of x_j appear in the expansion. In the physics literature, (1.1) usually is given in disguise as

$$Y(u, z_1)Y(v, z_2) \sim \sum_{n=0}^N Y(u_n v, z_2)(z_1 - z_2)^{-n-1},$$

with some $N \in \mathbb{N}$ depending on u and v , where the iteration of two vertex operators has been expanded, only the terms singular in $(z_1 - z_2)$ are given, and this relation should be implicitly understood to be used inside a correlation function.

The operator product expansion (1.1) is referred to as meromorphic since vertex operators acting on modules only involve integer powers of formal variables. But much of the interesting information in both physical and mathematical theories is hidden in the way different modules “interact”. So instead of vertex operators that always act from the underlying vertex operator algebra on a given module, *intertwining operators* have to be considered, which mediate between three (possibly distinct) modules and do not necessarily depend meromorphically on their variables.

While the existence and associativity of the operator product expansion for vertex operators $Y(\cdot, x) \cdot$ follows rather easily from the axioms, this is not the case for intertwining operators $\mathcal{Y}(\cdot, x) \cdot$. In fact, it is a deep result of the intricate $P(z)$ -tensor product theory of Huang and Lepowsky developed in [HL1]–[HL3] and [H1]. In [HLZ], Huang, Lepowsky and Zhang generalized their results such that they could drop the condition of rationality and of semisimplicity of the action of L_0 . In this way, a logarithmic dependence on the variables may appear in intertwining operators and correlation functions. This also seems to be the most successful rigorous treatment of logarithmic conformal field theory (e.g., see the review articles [G] and [F3]) so far.

Now we recall some of the results of [HLZ] which are central for this work. For any generalized module W of a vertex operator algebra V , its restricted dual $W' = \coprod_{n \in \mathbb{Z}} (W_{[n]})^*$ can be endowed with a module structure, where the associated vertex operator $Y'(\cdot, x) \cdot$ is defined by

$$\langle Y'(v, x)w', w \rangle = \langle w', Y(e^{xL_1}(-x^{-2})^{L_0}v, x^{-1})w \rangle$$

for all $v \in V$, $w \in W$ and $w' \in W'$. The pair (W', Y') is called the contragredient module to (W, Y) , and the contravariant functor $(\cdot)': (W, Y) \mapsto (W', Y')$ is called the contragredient functor. (Here and in the following, we understand a generalized module in the way a module is usually defined as for example in [LL], except that the action of L_0 may have a nilpotent part, i.e. the homogeneous subspaces $W_{[n]} = \{w \in W \mid (L_0 - n)^m w = 0 \text{ for } m \gg 0\}$ are generalized eigenspaces which are in particular assumed to be finite-dimensional. Sometimes we call such a structure simply a module, omitting the attribute generalized.) Also, let \mathcal{C} denote an as yet unspecified full subcategory of the category whose objects are all (generalized) modules of a given vertex operator algebra V , that is closed under the contragredient functor.

In [HLZ], the authors use the $P(z)$ -tensor product theory and carefully establish several conditions for the existence and associativity of the operator product expansion for logarithmic intertwining operators. Instead of repeating all the steps in their argument here, we refer to their paper for details and give only a summary of the results concerning the operator product expansion. Indeed, once an adequate subcategory \mathcal{C} has been chosen, it follows from Proposition 5.13 and Theorems 6.1 to 6.3 in [HLZ] that if its objects satisfy the following three conditions, then the nonmeromorphic operator product expansion exists and is associative. These conditions are:

- (i) All generalized V -modules in $\text{ob } \mathcal{C}$ are C_1 -cofinite, i.e. for all $W \in \text{ob } \mathcal{C}$ the space $W/C_1(W)$ is finite-dimensional with

$$C_1(W) = \text{span} \left\{ u_{-1}w \mid u \in \coprod_{n>0} V_{(n)}, w \in W \right\} .$$

- (ii) All generalized V -modules in $\text{ob } \mathcal{C}$ are quasi-finite-dimensional, i.e. for all $W \in \text{ob } \mathcal{C}$,

$$\dim \coprod_{n < N} W_{[n]} < \infty \quad \text{for all } N \in \mathbb{R} .$$

- (iii) Every object which is a finitely generated lower-truncated generalized V -module, except that it may have infinite-dimensional homogeneous subspaces, is an object in \mathcal{C} .

Note that the basic structure in [HLZ] are conformal vertex algebras, which are the same as vertex operator algebras except that they might not satisfy the lower-truncation condition and their homogeneous subspaces may be infinite-dimensional, and the same is true for their (generalized) modules. This is why the objects may not be assumed to have finite-dimensional homogeneous subspaces in condition (iii), but this must be proven.

The precise statement of the assertion is that under the above conditions, for any W_1, W_2, W_3, W'_4 and M in $\text{ob } \mathcal{C}$, any $P(z_1)$ -intertwining map I_1 of type $\binom{W_4}{W_1 M}$ and any $P(z_2)$ -intertwining map I_2 of type $\binom{M}{W_2 W_3}$, there is a $P(z_2)$ -intertwining map I of type $\binom{W_4}{W_1 \boxtimes_{P(z_0)} W_2 W_3}$ such that

$$\langle w'_4, I_1(w_1, z_1)I_2(w_2, z_2)w_3 \rangle = \langle w'_4, I(w_1 \boxtimes_{P(z_0)} w_2, z_2)w_3 \rangle \quad (1.2)$$

for all $w_1 \in W_1, w_2 \in W_2, w_3 \in W_3$ and $w'_4 \in W'_4$.

The formal similarity between (1.1) and (1.2) is obvious. The subtle point is that Huang, Lepowsky and Zhang really proved that in the operator product expansion of two logarithmic intertwining maps, there only appear powers of the variables and their logarithms (with no further dependence on the variables), while in the physics literature this is usually assumed without proof. With this information one can then try to find differential equations that are solved by the matrix elements of the product of two logarithmic intertwining maps at, say, z_1 and z_2 . (Such differential equations always do exist by a theorem of Huang [H2], [HLZ].) Expanding the solution in $z_0 = z_1 - z_2$ and z_2 , one arrives at the desired operator product expansion, evaluated inside a matrix element – and only this way is the operator product expansion well-defined.

In the present paper, we show that a certain family $\{\mathcal{W}(2, (2p-1)^{\times 3})\}_{p \geq 2}$ of \mathcal{W} -algebras satisfies the above conditions. The vertex operator algebra $\mathcal{W}(2, 3^{\times 3})$ is historically the first example of a “rational” *logarithmic* conformal field theory in the sense of [GK], i.e. a finite set of its modules closes under fusion. From the explicitly known characters for the vertex operator algebras $\mathcal{W}(2, (2p-1)^{\times 3})$ it follows that they all display logarithmic features, see [F2]. This is one of the motivations for us to study these \mathcal{W} -algebras in more detail here.

The main work of our proof is to establish that each member of this family is C_2 -cofinite, and from this fact condition (i) follows as we will show (the other two conditions are easy to check). Thus, we do not only prove the existence and associativity of the operator product expansion as described above for each $\mathcal{W}(2, (2p-1)^{\times 3})$, but we also establish one of the most useful and interesting properties in the study of vertex operator algebras for these \mathcal{W} -algebras.

Indeed, the condition of C_2 -cofiniteness was introduced by Zhu in [Z] and subsequently used to prove the convergence and modular invariance of characters for certain vertex operator algebras, and it is also related to his famous associative algebra $A(V)$. But C_2 -cofiniteness is also important because of its close

relation to rationality and regularity. It was proven by Li in [L] that any regular vertex operator algebra is also C_2 -cofinite, and Abe, Buhl and Dong were able to show in [ABD] that regularity is equivalent to rationality (in the sense of complete reducibility of all modules) and C_2 -cofiniteness together for vertex operator algebras of the form $V = \coprod_{m \in \mathbb{N}} V_{(m)}$ with $V_{(0)} = \mathbb{C}\Omega$. Even a conjecture was formulated that rationality (in the sense of complete reducibility of all modules), regularity and C_2 -cofiniteness are equivalent properties for vertex operator algebras, but $\mathcal{W}(2, 3^{\times 3})$ actually serves as a counter-example. This was known from the work of Kausch, and Abe explicitly noted it in [A]. Nevertheless, one may still conjecture the equivalence of C_2 -cofiniteness and “rationality” in the sense of [GK], i.e. a finite set of modules closes under fusion.

Our proof that all triplet algebras $V_{2p-1} = \mathcal{W}(2, (2p-1)^{\times 3})$ are C_2 -cofinite adds credibility to this conjecture. Indeed, from the C_2 -cofiniteness of V_{2p-1} it follows that the Zhu algebra $A(V_{2p-1})$ is finite-dimensional, and because of this there are only finitely many equivalence classes of indecomposable $A(V_{2p-1})$ -modules. This together with the strong restrictions coming from the structure of \mathcal{W} -algebras suggests that the assumedly equivalent properties both hold for all triplet algebras.

What makes this relationship particularly interesting is the fact that while rationality explicitly concerns the modules for a vertex operator algebra, the C_2 -cofiniteness condition can be studied solely in terms of the vertex operator algebra itself, without reference to any modules.

The remaining structure of the present paper is as follows. In section 2 we briefly recall a few basic properties concerning C_n -cofiniteness; we also explain what we mean by a \mathcal{W} -algebra in general and give some useful results. For standard notions concerning vertex operator algebras and for details about concepts like (logarithmic) intertwining maps (which are in one-to-one correspondence to (logarithmic) intertwining operators) or the $P(z)$ -tensor product, we refer the reader to the above-mentioned literature. Then, in section 3, we first prove our result for the intimately known vertex operator algebra $\mathcal{W}(2, 3^{\times 3})$ and then show how the respective arguments can be generalized to all members of the family $\{\mathcal{W}(2, (2p-1)^{\times 3})\}_{p \geq 2}$.

Acknowledgments: Nils Carqueville thanks Yi-Zhi Huang for explaining several aspects of $P(z)$ -tensor product theory to him. He also thanks Julia Voelskow for taking interest in this work. Both authors thank James Lepowsky and Geoffrey Buhl for many valuable comments. The research of Michael Flohr is supported by the European Union Network HPRN-CT-2002-00325 (EUCLID).

2 Preliminaries

The subspaces $C_n(W)$

In order to prove that the above conditions are satisfied for a given vertex operator algebra V and appropriately chosen \mathcal{C} , we first give two general properties of the spaces $C_n(W)$, where W is any generalized V -module. For $n \geq 2$, $C_n(W)$ is defined as $C_n(W) = \text{span}\{u_{-n}w \mid u \in V, w \in W\}$, and for $n = 1$, we have $C_1(W) = \text{span}\{u_{-1}w \mid u \in \coprod_{n>0} V_{(n)}, w \in W\}$. The space W is called C_n -cofinite if $\dim(W/C_n(W)) < \infty$. Thus, because of the L_{-1} -derivative property

$$Y(L_{-1}^m v, x) = \frac{d^m}{dx^m} Y(v, x),$$

it directly follows by comparing coefficients that

$$v_{-m-1} = \frac{1}{m} (L_{-1} v)_{-m} = \frac{1}{m!} (L_{-1}^m v)_{-1} \quad \text{for all } m \in \mathbb{Z}_+, \quad (2.1)$$

and hence every C_m -cofinite generalized V -module W is also C_n -cofinite for all $m \geq n \geq 1$. For $n = 1$, this can also be expressed by writing

$$C_1(W) = \text{span} \left\{ u_{-m} w \mid u \in \prod_{n>0} V_{(n)}, w \in W, m \in \mathbb{Z}_+ \right\}. \quad (2.2)$$

Secondly, $C_n(W)$ is invariant under the action of v_m for all $v \in V$ and $m \leq 0$. To prove this, one only needs to look at the well-known commutation relation

$$v_m u_{-n} w = u_{-n} v_m w + \sum_{i \in \mathbb{N}} \binom{m}{i} (v_i u)_{m-n-i} w \quad (2.3)$$

which follows from the Jacobi identity by performing the usual residue operation. For $m \leq 0$ the right-hand side of (2.3) obviously is an element of $C_n(W)$ because of the relation (2.1), and so this must also be true for the left-hand side. The result is

$$v_m C_n(W) \subset C_n(W) \quad \text{for all } v \in V, m \in \mathbb{Z}_{\leq 0}, n \in \mathbb{Z}_+. \quad (2.4)$$

Finally, another useful relation that follows from the Jacobi identity is

$$(u_m v)_n = \sum_{i \in \mathbb{N}} (-1)^i \binom{m}{i} u_{m-i} v_{n+i} - \sum_{i \in \mathbb{N}} (-1)^{i+m} \binom{m}{i} v_{m+n-i} u_i. \quad (2.5)$$

\mathcal{W} -algebras

A \mathcal{W} -algebra of type $\mathcal{W}(2, h_1, \dots, h_m)$ is a vertex operator algebra which has a generating set consisting of the vacuum Ω , the conformal vector ω of weight 2 and m additional primary vectors W^i of weight h_i , $i \in \{1, \dots, m\}$. The vertex operators or fields associated to these vectors are simple in the sense that they are not normal-ordered products of other fields. Sometimes the term \mathcal{W} -algebra is also used to refer to the algebra of modes instead of the vertex operator algebra.

What will be of particular importance for us is the notion of quasi-primary normal-ordered products which is due to Nahm [N]. When working with \mathcal{W} -algebras, we will mainly adopt his notation. For a more detailed exposition, see for example [F1]. Here, only those relations are given that are needed for our present purpose.

The usual normal-ordered product $:\phi_i(x)\phi_j(y): = \phi_i(x)_+\phi_j(y) + \phi_j(y)\phi_i(x)_-$ of two quasi-primary fields ϕ_i and ϕ_j is not necessarily quasi-primary for $x = y$. One of Nahm's results is that it is always possible to add certain correction terms, yielding a quasi-primary normal-ordered product denoted by $\mathcal{N}(\cdot, \cdot)$:

$$\begin{aligned} \mathcal{N}(\phi_j, \partial^n \phi_i) &= \sum_{r=0}^n (-1)^r \binom{n}{r} \binom{2(h_i + h_j + n - 1)}{r}^{-1} \binom{2h_i + n - 1}{r} \\ &\quad \cdot \partial^r N^{(h_i + n + r)}(\phi_j, \partial^{n-r} \phi_i) \\ &\quad - (-1)^n \sum_{\{k | h(ijk) \geq 1\}} C_{ij}^k \binom{h(ijk) + n - 1}{n} \\ &\quad \cdot \binom{2(h_i + h_j + n - 1)}{n}^{-1} \binom{2h_i + n - 1}{h(ijk) + n} \binom{\sigma(ijk) - 1}{h(ijk) - 1}^{-1} \\ &\quad \cdot \frac{\partial^{h(ijk) + n} \phi_k}{(\sigma(ijk) + n)(h(ijk) - 1)}. \end{aligned} \tag{2.6}$$

Here, $\{\phi_k\}_k$ is the family of quasi-primary fields of the corresponding \mathcal{W} -algebra, h_k are their respective weights, $h(ijk) := h_i + h_j - h_k$ and $\sigma(ijk) := h_i + h_j + h_k - 1$. The structure constants C_{ij}^k are defined such that $\sum_l C_{ij}^l d_{lk} = C_{ijk}$ with

$$C_{ijk} = \langle \Omega', (\phi_k)_{+h_k} (\phi_i)_{-h_k + h_j} (\phi_j)_{-h_j} \Omega \rangle \quad \text{and} \quad d_{ij} = \langle \Omega', (\phi_i)_{+h_i} (\phi_j)_{-h_j} \Omega \rangle,$$

and the $N^{(\cdot)}$ -product is defined by the relations

$$\begin{aligned} N^{(m)}(\phi, \psi)(x) &= \sum_{n \in \mathbb{Z}} x^{-n - h_\phi - h_\psi} N^{(m)}(\phi, \psi)_n, \\ N^{(m)}(\phi, \psi)_n &= \sum_{k < m} \phi_{n+k} \psi_{-k} + \sum_{k \geq m} \psi_{-k} \phi_{n+k} \end{aligned} \tag{2.7}$$

for any $m \in \mathbb{Z}$. The quasi-primary normal-ordered product of more than two fields is defined recursively, for example $\mathcal{N}(\phi_i, \phi_j, \phi_k) = \mathcal{N}(\phi_i, \mathcal{N}(\phi_j, \phi_k))$. If the

product of a field with itself is considered the notation is simplified, for example $\mathcal{N}(\psi, \psi) = \mathcal{N}(\psi^2)$.

Furthermore, in this notation the commutators of modes are given by

$$[(\phi_i)_m, (\phi_j)_n] = d_{ij} \delta_{m+n,0} \binom{h_i + m - 1}{2h_i - 1} + \sum_{\{k | h^{(ijk)} \geq 1\}} C_{ij}^k p_{h_i, h_j, h_k}(m, n) (\phi_k)_{m+n} \quad (2.8)$$

in terms of the polynomials

$$p_{h_i, h_j, h_k}(m, n) = \sum_{r, s \in \mathbb{N}} \delta_{r+s, h^{(ijk)}-1} a_{ijk}^r \binom{m+n-h_k}{r} \binom{h_i-n-1}{s}$$

with

$$a_{ijk}^r = \binom{2h_k+r-1}{r}^{-1} \binom{h_i+h_k-h_j+r-1}{r}.$$

In the next section, also the formal power series known as the character

$$\chi_V(q) = \text{tr}_V q^{L_0 - c/24} = q^{-c/24} \sum_{n \in \mathbb{N}} \dim V_{(n)} q^n$$

of the vertex operator algebra $V = \mathcal{W}(2, h_1, \dots, h_m)$ will be useful. In our proof of the existence of certain singular vectors, we will compare this character with the character of the vacuum Verma module of the \mathcal{W} -algebra. This is the induced module

$$U(\mathcal{W}(2, h_1, \dots, h_m)) \otimes_{U(\mathcal{W}(2, h_1, \dots, h_m)_{(+)})} \mathbb{C}_c,$$

where $U(\cdot)$ denotes the universal enveloping algebra of the \mathcal{W} -algebra, the space $\mathcal{W}(2, h_1, \dots, h_m)_{(+)}$ is defined by

$$\mathcal{W}(2, h_1, \dots, h_m)_{(+)} = \prod_{n \leq -1} \mathbb{C} L_{-n} \oplus \prod_{i=1}^m \prod_{n_i \leq h_i - 1} \mathbb{C} W_{-n_i}^i,$$

and \mathbb{C}_c is the trivial $\mathcal{W}(2, h_1, \dots, h_m)_{(+)}$ -module of central charge c . In other words, the vacuum Verma module is generated freely by the action of the modes L_n and W_n^i on a nonzero element Ω in \mathbb{C}_c , subject to the restrictions

$$L_n \Omega = 0 \text{ for all } n \geq -1 \quad \text{and} \quad W_n^i \Omega = 0 \text{ for all } n \geq -h_i + 1. \quad (2.9)$$

Because of these restrictions, the dimensions of the homogeneous subspaces $V_{(n)}$ are smaller than $p(n)$, where $p(n)$ is the number of partitions of n into sums of positive integers, generated by the function

$$(\varphi(q))^{-1} = \prod_{n \geq 1} (1 - q^n)^{-1} = \sum_{n \in \mathbb{N}} p(n) q^n.$$

Taking the restrictions (2.9) into account, the vacuum Verma module character is given by

$$\chi_V^{\text{Verma}}(q) = q^{-c/24}(\varphi_2(q))^{-1} \prod_{i=1}^m (\varphi_{h_i}(q))^{-1}, \quad (2.10)$$

where we have introduced the generating functions φ_k , $k \geq 2$, as truncated φ -functions:

$$\varphi_k(q) = \prod_{n \geq k} (1 - q^n) = \varphi(q) \prod_{l=1}^{k-1} (1 - q^l)^{-1}. \quad (2.11)$$

3 Properties of the triplet algebras

The triplet algebra at $c = -2$

With the general relations of the preceding section at hand, we now choose a vertex operator algebra V and a suitable category \mathcal{C} for which we will prove the conditions for the existence and associativity of the nonmeromorphic operator product expansion. We take $V = V_3$ which is the triplet algebra with central charge $c = -2$ [Kau], [F2], [GK] and \mathcal{C} such that its objects are exactly all finitely generated lower-truncated generalized V -modules. (Note that in this context, the notion of such a module of V by definition encompasses a possible Jordan cell structure in the L_0 -grading.) In particular, this choice includes all (generalized) highest weight modules, but also those on which L_1 acts only nilpotently (and not necessarily trivially) on the generating vector.

The triplet algebra at $c = -2$ is a \mathcal{W} -algebra of type $\mathcal{W}(2, 3^{\times 3})$. It is generated by the modes L_m of the Virasoro field $T(x) = \sum_{m \in \mathbb{Z}} L_m x^{-m-2}$ associated to the vector ω of weight 2 which implements the conformal symmetry, and the modes W_m^a of a triplet (under the action of the group $SO(3)$) of primary fields of weight 3, $W^a(x) = \sum_{m \in \mathbb{Z}} W_m^a x^{-m-3}$ with $a \in \{\pm 1, 0\}$, which “maximally extend” the conformal symmetry, acting on some module $W \in \text{ob } \mathcal{C}$. With this notation and by relation (2.2), the vectors

$$L_{-m+1}w \text{ and } W_{-m}^a w \text{ are in } C_1(W) \text{ for all } w \in W \text{ and all } m \geq 3, \quad (3.1)$$

while for all other values of m , this is not necessarily the case.

By this choice of \mathcal{C} , condition (iii) above is satisfied. The fact that the homogeneous subspaces of the (generalized) modules in $\text{ob } \mathcal{C}$ are really finite-dimensional follows from results of Buhl [B] on a module spanning set, using the fact that they are finitely generated and all triplet algebras are C_2 -cofinite, see below.

Condition (ii) also is satisfied: by the action of any mode v_n with $v \in V$, the weight of an element to which v_n is applied to changes by an integer value, and there are only finitely many vectors that generate V , namely ω and W^a (together with the vacuum Ω).

In order to see that condition (i) is satisfied as well, i.e. $C_1(W)$ is finite-codimensional for all $W \in \text{ob } \mathcal{C}$, we assume without loss of generality that W is generated by some element $w = w^{(0)}$ together with its (finitely many) “logarithmic partners” $w^{(i)}$. Then by (2.3) and (2.5), every vector in W is a linear combination of elements of the form

$$\mathcal{M}_{-m_1} \dots \mathcal{M}_{-m_k} L_{-1}^M \prod_{a \in \{\pm 1, 0\}} \left((W_{-2}^a)^{N_2^a} (W_{-1}^a)^{N_1^a} (W_0^a)^{N_0^a} \right) \mathcal{M}_{n_1} \dots \mathcal{M}_{n_l} w^{(i)} \quad (3.2)$$

where \mathcal{M} is a placeholder for either L or W^a ; $M, N_0^a, N_1^a, N_2^a \in \mathbb{N}$, $n_1, \dots, n_l \in \mathbb{Z}_+$ and $m_1, \dots, m_k \in \mathbb{Z}_{\geq 2}$ for $\mathcal{M} = L$ while $m_1, \dots, m_k \in \mathbb{Z}_{\geq 3}$ for $\mathcal{M} = W^a$.

In the case that k is strictly larger than zero, (3.1) immediately shows that any such element is in $C_1(W)$. On the other hand, for $k = 0$ there are only finitely many possibilities for the term $\mathcal{M}_{n_1} \dots \mathcal{M}_{n_l} w^{(i)}$ not to vanish because of the lower-truncatedness of W . The factor $(W_0^a)^{N_0^a}$ can also do no harm as it does not change the generalized weight of the element it is applied to, and each Jordan cell is finite-dimensional by the definition of \mathcal{C} , given the fact that condition (ii) is satisfied.

So what deserves special attention are the powers of L_{-1} , W_{-1}^a and W_{-2}^a in the case $k = 0$, because when applied to some element of W , the result need not be in $C_1(W)$, but each of these modes strictly increases the generalized weight. As there is certainly no “upper-truncation condition” for the module W , the appearance of these modes in (3.2) makes it seem possible that the complement of $C_1(W)$ in W is infinite-dimensional.

But fortunately, in this situation a theorem due to Buhl (see [B], Theorem 1) applies, which is a generalization of an earlier result of Gaberdiel and Neitzke [GN]. It states the following (among other things): If a vertex operator algebra V is C_2 -cofinite, i.e. $\dim(V/C_2(V)) < \infty$ with $C_2(V) = \text{span}\{u_{-2}v \mid u, v \in V\}$, then every weak V -module W is spanned by elements of the form

$$x_{-n_1}^1 \dots x_{-n_k}^k w \quad (3.3)$$

with $w \in W$, $n_1 \geq \dots \geq n_k > -L$, where L is some fixed number, and the vectors $x^1, \dots, x^k \in V$ are representatives of the elements of a basis of $V/C_2(V)$. In addition, if $n_j \leq 0$, then $n_i = n_j$ for at most Q indices i , where Q is another fixed number. This last feature is the most important one for the present situation as it implies that only a limited number of powers of L_{-1} , W_{-1}^a and W_{-2}^a has to be considered if V is C_2 -cofinite.

(Note that in (3.3) the convention for the indices of modes is the one used most often in the mathematics literature, i.e. any vertex operator is expanded into a series $\sum_{n \in \mathbb{Z}} v_n x^{-n-1}$ regardless of the weight of the associated vector v . On the other hand, in the physics literature it is common to expand a field that is associated to a vector u of weight h into a series $\sum_{n \in \mathbb{Z}} u_n^{\text{phys}} x^{-n-h}$. The latter

convention is used here only in the context of \mathcal{W} -algebras. When comparing results expressed in differing notations, the relation $u_n = u_{n-h+1}^{\text{phys}}$ is used.)

The triplet algebra at $c = -2$ has the virtue of being C_2 -cofinite because of the existence of certain singular vectors. Several authors (see [GN] and [M]) have been aware of this fact for some time, and it was recently proven by Abe in [A]. The proof given here uses a different method. Indeed, the explicit form of six singular vectors at level 6 is known [GK], [R]:

$$\begin{aligned} N^{ab} = & W_{-3}^a W_{-3}^b \Omega - \delta_{ab} \left(\frac{8}{9} L_{-2}^3 + \frac{19}{36} L_{-3}^2 + \frac{14}{9} L_{-4} L_{-2} - \frac{16}{9} L_{-6} \right) \Omega \\ & + i \varepsilon_{abc} \left(-2 W_{-4}^c L_{-2} + \frac{5}{4} W_{-6}^c \right) \Omega. \end{aligned} \quad (3.4)$$

In order to prove that V is really C_2 -cofinite, we first observe that in the above expression for the singular vector N^{ab} , because of (2.1) each term that it is made of is manifestly in $C_2(V)$ except for $W_{-3}^a W_{-3}^b \Omega$ and $L_{-2}^3 \Omega$. As any singular vector is divided out in the vertex operator algebra of interest, it follows that for $a \neq b$,

$$W_{-3}^a W_{-3}^b \Omega \in C_2(V) \quad \text{and} \quad \left((W_{-3}^a)^2 - (W_{-3}^b)^2 \right) \Omega \in C_2(V).$$

Since W_{-3}^a leaves the space $C_2(V)$ invariant (recall (2.4)), $W_{-3}^a ((W_{-3}^a)^2 - (W_{-3}^b)^2) \Omega$ is an element of $C_2(V)$ as well. But this element can also be written as

$$(W_{-3}^a)^3 \Omega - W_{-3}^a (W_{-3}^b)^2 \Omega = (W_{-3}^a)^3 \Omega - W_{-3}^b W_{-3}^a W_{-3}^b \Omega + Y_{-6} W_{-3}^b \Omega, \quad (3.5)$$

where $Y_{-6} = [W_{-3}^a, W_{-3}^b]$ applied to any vector $v \in V$ yields an element of $C_2(V)$ because in the commutator of modes of primary fields of weight 3 there can only appear modes corresponding to fields of weight less than or equal to 5 (see equation (2.8) above). So in particular, the last term in (3.5) is in $C_2(V)$. In addition, the second last term in this equation also is in $C_2(V)$ as W_{-3}^b leaves this space invariant. Hence, it follows that $(W_{-3}^a)^3 \Omega \in C_2(V)$, and we have

$$(W_{-3}^a)^m \Omega \in C_2(V) \quad \text{for all } m \geq 3. \quad (3.6)$$

From this and the fact that $((W_{-3}^a)^2 - \frac{8}{9} L_{-2}^3) \Omega$ is in $C_2(V)$ it follows that $(W_{-3}^a)^2 L_{-2}^3 \Omega \in C_2(V)$. Now using the invariance of $C_2(V)$ under L_{-2} and W_{-3}^a one more time it is easy to see that

$$\left((W_{-3}^a)^2 - \frac{8}{9} L_{-2}^3 \right)^2 \Omega = \left((W_{-3}^a)^4 + \frac{64}{81} L_{-2}^6 - \frac{8}{9} (W_{-3}^a)^2 L_{-2}^3 - \frac{8}{9} L_{-2}^3 (W_{-3}^a)^2 \right) \Omega$$

is an element of $C_2(V)$. But from the above discussion it is also clear that each term on the right-hand side apart from $\frac{64}{81} L_{-2}^6 \Omega$ is in $C_2(V)$, and so it follows that $L_{-2}^6 \Omega$ must be an element of $C_2(V)$ as well.

As a consequence of the reasoning of the last paragraph, sufficiently large powers of L_{-2} and W_{-3}^a (6 or maybe less in the first case, 3 or maybe less in the latter) applied to any element in V yield elements in $C_2(V)$. Thus it is proven that $C_2(V)$ is finite-codimensional.

Now that it has been shown that the prerequisite of Buhl's theorem is satisfied for the triplet algebra at $c = -2$, it can be used since by definition any object in \mathcal{C} is a weak module and the elements in (3.2) are of the same form as those in (3.3). This means that if it can be argued that ω and W^a are *not* in $C_2(V)$ and can thus be taken to be representatives of elements in a basis for $V/C_2(V)$, there actually *is* some sort of an "upper-truncation condition", but for the exponents of the modes L_{-1} , W_{-1}^a and W_{-2}^a in (3.2). So for $k = 0$, it follows that only finitely many elements of the form (3.2) span the " $(k = 0)$ -part" of W . This is exactly the statement that W is C_1 -cofinite.

It remains to be seen that ω and W^a are not in $C_2(V)$. For the moment, consider the possibility that ω *is* in $C_2(V)$. Then there must be $u, v \in V$ such that $u_{-2}v = \omega$. By comparing weights on both sides, we arrive at the condition $\text{wt } u + \text{wt } v + 1 = 2$. But since the vertex operator algebra V under consideration is of the form $V = \coprod_{n \in \mathbb{N}} V_{(n)}$ with $V_{(0)} = \mathbb{C}\Omega$, this condition says that either u or v must be (a scalar multiple of) the vacuum (and the other one of weight 1). This is not possible for the conformal vector, leading to a contradiction. By a similar reasoning, one also sees that $W^a \notin C_2(V)$.

Finally, it needs to be shown that the chosen category is closed with respect to the contragredient functor. By the definition of the graded dual $W' = \prod_{n \in \mathbb{Z}} (W_{[n]})^*$ it is clear that it is lower-truncated. In order to establish that it is also finitely generated, choose a minimal generating set $\{w_1, \dots, w_N\} \subset W \in \text{ob } \mathcal{C}$ from a basis $\bigcup_{n \in \mathbb{Z}} B_n$ of W , where B_n is a basis of $W_{[n]}$ for all $n \in \mathbb{Z}$. Then all $w \in W$ are linear combinations of elements of the form

$$\mathcal{M}_{n_1} \dots \mathcal{M}_{n_k} w_i ,$$

where \mathcal{M} denotes the same as in (3.2). Let w'_1, \dots, w'_N be the elements of the dual basis in W' such that $\langle w'_i, w_j \rangle = \delta_{ij}$. Because of this, all $w' \in W'$ that may give a nonvanishing matrix element with some $w \in W$ must be linear combinations of elements of the form $\mathcal{M}'_{n_k} \dots \mathcal{M}'_{n_1} w'_i$. To see this, assume that there is an element $\tilde{w}' \notin \{w'_1, \dots, w'_N\}$ in W' such that $\{\tilde{w}', w'_1, \dots, w'_N\}$ is a subset of a minimal set of generating vectors of W' . It follows that $\langle \tilde{w}', w_i \rangle = 0$ for all $i \in \{1, \dots, N\}$ and thus

$$\begin{aligned} \langle \mathcal{M}'_{-m_1} \dots \mathcal{M}'_{-m_k} \tilde{w}', \mathcal{M}_{-n_1} \dots \mathcal{M}_{-n_l} w_i \rangle &= \langle \tilde{w}', \mathcal{M}'_{m_k} \dots \mathcal{M}'_{m_1} \mathcal{M}_{-n_1} \dots \mathcal{M}_{-n_l} w_i \rangle \\ &= \delta_{\sum_i m_i, \sum_j n_j} \left\langle \tilde{w}', \sum_{\{I \mid \text{wt } w_I = \text{wt } w_i\}} a_I w_I \right\rangle \\ &= 0 , \end{aligned}$$

where $a_I \in \mathbb{C}$ are the coefficients that result from applying the commutation relations of the \mathcal{M} -modes. This means that the subspace of generalized weight $\text{wt } \tilde{w}'$ has a dimension that is strictly larger than the dimension of the corresponding subspace in W . But by the definition of the graded dual of W , these finite-dimensional subspaces must have the same dimension, so there cannot be an element \tilde{w}' as above, and \mathcal{C} is closed under the contragredient functor.

The triplet algebras at $c_{p,1}$

The triplet algebra at $c = -2$ is only the first member of an infinite family of triplet \mathcal{W} -algebras $\{\mathcal{W}(2, (2p-1)^{\times 3})\}_{p \geq 2}$ with central charge $c_{p,1} = 1 - 6(p-1)^2/p$, where for each $p \in \mathbb{Z}_{\geq 2}$ the three primary fields of weight $2p-1$ are a triplet under the action of the group $SO(3)$, which means that the structure constant $C_{W^a, W^b}^{W^c}$ is proportional to ε_{abc} [Kau]. It is the goal of this section to show that the above conditions (i), (ii) and (iii) are also satisfied in this general case.

If one defines the category \mathcal{C} analogously to the special case of $p = 2$, one immediately sees that the conditions of quasi-finite dimensionality and of finitely generated lower-truncated generalized modules in $\text{ob } \mathcal{C}$ hold in the same way as before with the obvious generalization of the arguments. What requires additional work is to establish the C_1 -cofiniteness of all objects in \mathcal{C} .

Let V_Δ denote the vertex operator algebra associated to the \mathcal{W} -algebra $\mathcal{W}(2, \Delta^{\times 3})$ for a fixed $\Delta := 2p-1$ with $p \in \mathbb{Z}_{\geq 3}$. If V_Δ is C_2 -cofinite, one can apply Buhl's theorem as in the case $p = 2$, and any V_Δ -module under consideration would be C_1 -cofinite, which together with the other properties of V_Δ gives the existence and associativity of the nonmeromorphic operator product expansion. Compared with the case $p = 2$, the difficulty of proving the C_2 -cofiniteness of V_Δ stems from the lack of explicit expressions for singular vectors that are crucial for a proof of C_2 -cofiniteness. A priori, it is not even clear whether such singular vectors at all exist for arbitrary $p \in \mathbb{Z}_{\geq 3}$.

As it turns out, one can argue for the existence of certain singular vectors of weight $2(2p-1)$ with the help of the explicitly known character of V_Δ that was obtained in [F2]. By analyzing this character in detail one can show that for arbitrary $p \in \mathbb{Z}_{\geq 2}$, singular vectors of the form

$$\begin{aligned} N^{ab} &= W_{-\Delta}^a W_{-\Delta}^b \Omega + \delta_{ab} (\text{Virasoro-polynomial}) \Omega \\ &\quad + \varepsilon_{abc} (\text{Virasoro-} W_m^c \text{-polynomial}) \Omega \end{aligned} \quad (3.7)$$

exist, where in the last term only summands with exactly one W^c -mode can appear (of course, equation (3.4) is of this form, too).

In order to show that singular vectors as in (3.7) really exist, we first recall the character

$$\chi_{V_\Delta}(q) = \frac{q^{-1/24}}{\varphi(q)} \sum_{n \in \mathbb{Z}} (2n+1) q^{(2np+p-1)^2/(4p)} \quad (3.8)$$

from [F2]. If we expand both this character and the vacuum Verma module character $\chi_{V_\Delta}^{\text{Verma}}(q)$ given by (2.10) into a formal power series in q and compare the coefficients of $q^{(2p-1)+3}$ (times $q^{-c_{p,1}/24}$), we see that the dimensions of the homogeneous subspaces of weight $(2p-1)+3$ of the vacuum Verma module and the \mathcal{W} -algebra itself differ by 3. The reason for this is the following: From the Kac determinant it follows that the Virasoro algebra of central charge $c_{p,1}$ has an infinite set of highest weight modules where the highest weights are given by $h_{2k-1,1} = (k-1)(kp-1)$, $k \in \mathbb{Z}_+$. By a standard argument it follows that these modules have singular vectors at level $2k-1$. In particular, for $k=2$ the highest weight vectors of weight $2p-1$ can be identified with the vectors $W_{-\Delta}^a \Omega$ as we have $\Delta = 2p-1$. So because of the additional structure of the \mathcal{W} -algebra with its fields W^a , pure Virasoro modules are embedded into the full vertex operator algebra $\mathcal{W}(2, \Delta^{\times 3})$, and the difference of the dimensions above is due to the three singular vectors of weight $(2p-1)+3$.

If these three vectors are divided out of the vacuum Verma module, we obtain a structure to which the character

$$\tilde{\chi}(q) = q^{-c_{p,1}/24} \left(\frac{1}{\varphi_2(q)} + \frac{3q^{2p-1}(1-q^3)}{\varphi(q)(\varphi_{2p-1}(q))^2} \right) \quad (3.9)$$

pertains, where we use the notation introduced in (2.11). The first term in this expression accounts for the action of the Virasoro algebra on the vacuum alone. The second term reflects the fact that beginning at level $2p-1$, the modes associated to the three distinct W^a -fields act nontrivially on the vacuum. With respect to the Virasoro algebra, this is a highest weight vector, which explains the factor $q^{2p-1}/\varphi(q)$. Furthermore, the factor $1-q^3$ is due to the singular vectors of weight $(2p-1)+3$ discussed above, and the term $(\varphi_{2p-1}(q))^{-2}$ comes from the action of the W^a -modes on the vacuum. The second power (and not the third) has to be taken here in order not to doubly count the contribution from the W^a -modes because of the three-fold multiplicity.

Partially expanding both (3.8) and (3.9) into a formal power series yields

$$\begin{aligned} \chi_{V_\Delta}(q) &= \frac{q^{-c_{p,1}/24}}{\varphi(q)} (1 - q + 3q^{2p-1} - 3q^{2p+2} + \mathcal{O}(q^{6p-2})) , \\ \tilde{\chi}(q) &= \frac{q^{-c_{p,1}/24}}{\varphi(q)} (1 - q + 3q^{2p-1} - 3q^{2p+2} + 6q^{4p-2} + \mathcal{O}(q^{4p-1})) . \end{aligned}$$

We are interested in the dimensions of the homogeneous subspaces of weight $2\Delta = 4p-2$ described by these characters. Comparing the coefficients of q^{4p-2} (times $q^{-c_{p,1}/24}$) by taking the relevant contributions from $(\varphi(q))^{-1} = \sum_{n \in \mathbb{N}} p(n)q^n$ into account, we see that these dimensions differ by 6. Thus we have found that six additional singular vectors of weight 2Δ are divided out in $\mathcal{W}(2, \Delta^{\times 3})$. The reason that these vectors must involve a term with two W^a -modes is that there

are no pure Virasoro singular vectors of weight 2Δ . Finally, the form of (3.7) is a direct consequence of the $SO(3)$ -structure of $\mathcal{W}(2, \Delta^{\times 3})$ [Kau].

We now continue the proof of C_2 -cofiniteness of the \mathcal{W} -algebras $\mathcal{W}(2, \Delta^{\times 3})$. As in the special case $p = 2$ it is clear that nearly all possible vectors in the expression (3.7) for the singular vector N^{ab} are elements of $C_2(V_\Delta)$ because of the fact that $W_m^a \Omega = 0$ for all $m \geq -\Delta + 1$. The only vectors for which this might not be true are $W_{-\Delta}^a W_{-\Delta}^b \Omega$ and $\alpha L_{-2}^\Delta \Omega$, the latter appearing in the δ_{aa} -term in N^{ab} . If it can be shown that the coefficient α is not zero, the exact same reasoning as in the case $p = 2$ can be applied to see that V_Δ is C_2 -cofinite. So the question that remains to be answered is whether or not $\alpha \neq 0$.

To find the correct answer, we first observe that the vertex operator to which a singular vector corresponds necessarily is a primary field. In particular, it is a quasi-primary field. As the vector $W_{-\Delta}^a W_{-\Delta}^b \Omega$ appears in the expression for the singular vector N^{ab} , the corresponding quasi-primary null-field must be a linear combination of quasi-primary fields, and one of these must be the normal-ordered product $\mathcal{N}(W^a, W^b)$.

The next step is to note that the quasi-primary field $\mathcal{N}(W^a, W^b)$ alone cannot be the null-field. To see this, we make use of the fact that the mode L_1 annihilates the vector N^{ab} . Thus, by expanding the null-field into modes,

$$\begin{aligned} L_1 N^{aa} &= L_1 (W_{-\Delta}^a W_{-\Delta}^a \Omega + (\text{Virasoro-polynomial}) \Omega) \\ &\stackrel{\Delta-1}{=} L_1 (W_{-\Delta}^a W_{-\Delta}^a \Omega + \beta L_{-4} L_{-2}^{\Delta-2} \Omega + \gamma L_{-3}^2 L_{-2}^{\Delta-3} \Omega) \\ &\stackrel{\Delta-1}{=} 0. \end{aligned} \tag{3.10}$$

Here, the symbol $\stackrel{\Delta-1}{=}$ has been introduced, which means “equal to, modulo vectors with less than $\Delta - 1$ modes applied to the vacuum Ω ”. For example,

$$\beta L_{-4} L_{-2}^{\Delta-2} \Omega + \gamma L_{-3}^2 L_{-2}^{\Delta-3} \Omega \stackrel{\Delta-1}{=} \beta L_{-4} L_{-2}^{\Delta-2} \Omega + \gamma L_{-3}^2 L_{-2}^{\Delta-3} \Omega + \delta L_{-4}^2 L_{-2}^{\Delta-4} \Omega.$$

So far, the values of the constants β and γ are unknown. If the null-field were equal to $\mathcal{N}(W^a, W^b)$, the coefficients $\beta = \beta_{WW}$ and $\gamma = \gamma_{WW}$ could be computed from the above formula (2.6) for quasi-primary normal-ordered products in terms of the structure constant $C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}$. In principle, this constant can be computed for each $p \in \mathbb{Z}_{\geq 3}$ separately, but neither are such computations carried out easily nor is it necessary to know the exact value of the constant; only the information that it is not zero is crucial.

With this, a straight-forward calculation using (3.10) shows that

$$L_2 (W_{-\Delta}^a W_{-\Delta}^a + \beta_{WW} L_{-4} L_{-2}^{\Delta-2} + \gamma_{WW} L_{-3}^2 L_{-2}^{\Delta-3}) \Omega \stackrel{\Delta-1}{\neq} 0.$$

So the field $\mathcal{N}(W^a, W^b)$ is quasi-primary but not primary and can thus not be the null-field. Instead, other quasi-primary fields must be added to $\mathcal{N}(W^a, W^b)$ to get the null-field. Of all these fields, only those are of immediate interest

that yield primarity of the null-field *at length* $\Delta - 1$, i.e. $L_2 N^{aa} \stackrel{\Delta-1}{=} 0$. Define \mathcal{X} to be the set of all quasi-primary fields of weight 2Δ except $\mathcal{N}(T^\Delta)$ in whose mode expansion appear Virasoro-monomials up to degree $\Delta - 1$; in particular, $L_{-4}L_{-2}^{\Delta-2}$ is such a monomial. For example, $\mathcal{N}(\partial^2 T, \mathcal{N}(T^{\Delta-2})) \in \mathcal{X}$. Then the singular vector associated to the null-field satisfies the identity

$$\begin{aligned} N^{aa} \stackrel{\Delta-1}{=} & \left((\mathcal{N}(W^a, W^a))_{-2\Delta} + \alpha (\mathcal{N}(T^\Delta))_{-2\Delta} + \sum_{X \in \mathcal{X}} k_X X_{-2\Delta} \right) \Omega \\ & \stackrel{\Delta-1}{=} \left(W_{-\Delta}^a W_{-\Delta}^a + \alpha L_{-2}^\Delta + (\beta_{T^\Delta} + \beta_{WW}) L_{-4} L_{-2}^{\Delta-2} \right. \\ & \quad \left. + (\gamma_{T^\Delta} + \gamma_{WW}) L_{-3}^2 L_{-2}^{\Delta-3} + \sum_{X \in \mathcal{X}} (\beta_X L_{-4} L_{-2}^{\Delta-2} + \gamma_X L_{-3}^2 L_{-2}^{\Delta-3}) \right) \Omega. \end{aligned}$$

(Note that there are no vectors of length $\Delta - 1$ in $L_2 L_{-3}^2 L_{-2}^{\Delta-3} \Omega$, so the γ -terms do not have to be considered when L_2 acts on N^{aa} .)

Now the assumption is made that $\alpha = 0$. Then one can use the fact that $L_2 N^{aa} = 0$ to find an explicit expression for the parameter $B := \sum_{X \in \mathcal{X}} \beta_X$ in terms of the structure constant $C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}$. (Fields \mathcal{F} of weight $2\Delta - 1$ like $\mathcal{N}(\partial T, T^{\Delta-2})$ with one derivative term need not be taken into account since the structure constants $C_{W^a, W^a}^{\mathcal{F}}$ for such fields vanish, see [BFKNRV].) To do this, we need to know in which exact way β_{WW} is proportional to $C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}$, so that we have $\beta_{WW} = \beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}$ with β'_{WW} a nonzero constant whose exact value can be calculated to be $\beta'_{WW} = -\frac{(2\Delta-1)(\Delta-1)}{2(4\Delta-3)}$ by equation (2.6). With this notation we arrive at

$$\begin{aligned} 0 = L_2 N^{aa} \stackrel{\Delta-1}{=} & L_2 \left(W_{-\Delta}^a W_{-\Delta}^a + \left(\beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + B \right) L_{-4} L_{-2}^{\Delta-2} \right) \Omega \\ & \stackrel{\Delta-1}{=} \left([L_2, W_{-\Delta}^a W_{-\Delta}^a] + 6 \left(\beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + B \right) L_{-2}^{\Delta-1} \right) \Omega \\ & \stackrel{\Delta-1}{=} \left([L_2, W_{-\Delta}^a] W_{-\Delta}^a + 6 \left(\beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + B \right) L_{-2}^{\Delta-1} \right) \Omega \\ & \stackrel{\Delta-1}{=} \left((2(\Delta-1) + \Delta) W_{2-\Delta}^a W_{-\Delta}^a + 6 \left(\beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + B \right) L_{-2}^{\Delta-1} \right) \Omega \\ & \stackrel{\Delta-1}{=} \left((3\Delta-2) [W_{2-\Delta}^a, W_{-\Delta}^a] + 6 \left(\beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + B \right) L_{-2}^{\Delta-1} \right) \Omega \\ & \stackrel{\Delta-1}{=} \left((3\Delta-2) C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + 6 \left(\beta'_{WW} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} + B \right) \right) L_{-2}^{\Delta-1} \Omega, \end{aligned}$$

where it has been used in the last line that $p_{\Delta, \Delta, 2\Delta-2}(2-\Delta, -\Delta) = 1$. The above equation holds if and only if

$$B = -\frac{6\Delta^2 - 8\Delta + 3}{6(4\Delta - 3)} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}. \quad (3.11)$$

The idea to prove that $\alpha \neq 0$ now is to find another way to explicitly compute the value of B that does *not* agree with the one given in (3.11). But before this is done it should be noticed that the parameters $\beta = \beta_{WW} + B$ and $\gamma = \gamma_{WW} + \sum_{X \in \mathcal{X}} \gamma_X$ can be expressed completely in terms of the structure constant $C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}$ and B : β_{WW} and γ_{WW} can be calculated by equation (2.6), and from the fact that each field X in \mathcal{X} is quasi-primary (which means $L_1 X_{-2\Delta} \Omega \stackrel{\Delta-1}{=} 0$ among other things) it follows that $\sum_{X \in \mathcal{X}} \gamma_X = -\frac{5}{8}B$. As a consequence we have

$$\beta = -\frac{1}{2} \frac{2\Delta - 1}{4\Delta - 3} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} (\Delta - 1) + B, \quad (3.12a)$$

$$\gamma = -\frac{1}{2} \frac{2\Delta - 1}{4\Delta - 3} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \left((\Delta - 2)^2 - \frac{1}{2}(\Delta - 2)(\Delta - 3) \right) - \frac{5}{8}B. \quad (3.12b)$$

These relations will be made use of without explicit mention in the following.

The vector $N_{-2\Delta}^{aa} \Omega$ is already completely known at length $\Delta - 1$ up to the structure constant $C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}$, and the same situation will now be achieved for the vector $N_{-2\Delta-1}^{aa} \Omega$ as an intermediate step. For this, the relation $[L_m, \phi_n] = ((h-1)m - n)\phi_{m+n}$ with $m \in \{\pm 1, 0\}$ for a quasi-primary field ϕ of weight h is employed: at length $\Delta - 1$ we see that $[L_{-1}, N_{-2\Delta}^{aa}] \Omega = L_{-1} N^{aa}$ is equal to

$$\begin{aligned} & L_{-1} (W_{-\Delta}^a W_{-\Delta}^a + \beta L_{-4} L_{-2}^{\Delta-2} + \gamma L_{-3}^2 L_{-2}^{\Delta-3}) \Omega \\ \stackrel{\Delta-1}{=} & (W_{-\Delta}^a [L_{-1}, W_{-\Delta}^a] + [L_{-1}, W_{-\Delta}^a] W_{-\Delta}^a + 3\beta L_{-5} L_{-2}^{\Delta-2} \\ & + (\Delta - 2)\beta L_{-4} L_{-3} L_{-2}^{\Delta-3} + 4\gamma L_{-4} L_{-3} L_{-2}^{\Delta-3} + (\Delta - 3)\gamma L_{-3}^3 L_{-2}^{\Delta-4}) \Omega \\ \stackrel{\Delta-1}{=} & (2W_{-\Delta-1}^a W_{-\Delta}^a + [W_{-\Delta}^a, W_{-\Delta-1}^a]) \\ & + 3\beta L_{-5} L_{-2}^{\Delta-2} + ((\Delta - 2)\beta + 4\gamma) L_{-4} L_{-3} L_{-2}^{\Delta-3} + (\Delta - 3)\gamma L_{-3}^3 L_{-2}^{\Delta-4}) \Omega \\ \stackrel{\Delta-1}{=} & C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} p_{\Delta, \Delta, 2\Delta-2}(-\Delta, -\Delta - 1) \left((\Delta - 1)L_{-5} L_{-2}^{\Delta-2} \right. \\ & \left. + (\Delta - 1)(\Delta - 2)L_{-4} L_{-3} L_{-2}^{\Delta-3} + \binom{\Delta - 1}{3} L_{-3}^3 L_{-2}^{\Delta-4} \right) \Omega \\ & + (3\beta L_{-5} L_{-2}^{\Delta-2} + ((\Delta - 2)\beta + 4\gamma) L_{-4} L_{-3} L_{-2}^{\Delta-3}) \Omega \\ & + (\Delta - 3)\gamma L_{-3}^3 L_{-2}^{\Delta-4} \Omega. \end{aligned} \quad (3.13)$$

But because of the quasi-primarity of the vector N^{aa} , this must also be equal to $N_{-2\Delta-1}^{aa} \Omega$. Of course the latter is not known explicitly, but at length $\Delta - 1$ the relevant parameters can be inferred. Firstly, there is a contribution to $N_{-2\Delta-1}^{aa} \Omega$ from $\mathcal{N}(W^a, W^a)_{-2\Delta-1} \Omega$, and only the terms of length $\Delta - 1$ will be of importance here. Secondly, the contribution of the fields in \mathcal{X} has to be taken into account. Computing this contribution exactly would require the knowledge of the exact values of the parameters k_X in

$$N^{aa} \stackrel{\Delta-1}{=} \mathcal{N}(W^a, W^a)_{-2\Delta} \Omega + \sum_{X \in \mathcal{X}} k_X X_{-2\Delta} \Omega.$$

These are not available, but all we really need to know in this case are the coefficients of the relevant monomials at length $\Delta - 1$. Denoting these coefficients by ξ_i , $i \in \{1, 2, 3\}$, (3.13) is also equal to

$$\begin{aligned}
& \mathcal{N}(W^a, W^a)_{-2\Delta-1}\Omega + (\xi_1 L_{-5} L_{-2}^{\Delta-2} + \xi_2 L_{-4} L_{-3} L_{-2}^{\Delta-3} + \xi_3 L_{-3}^3 L_{-2}^{\Delta-4}) \Omega \\
\stackrel{\Delta-1}{=} & \frac{1}{4} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \frac{2\Delta-1}{4\Delta-3} (-n-2\Delta+2)(-n-2\Delta+1) \Big|_{n=-2\Delta-1} \\
& \cdot \left((\Delta-1) L_{-5} L_{-2}^{\Delta-2} + (\Delta-1)(\Delta-2) L_{-4} L_{-3} L_{-2}^{\Delta-3} \right. \\
& \quad \left. + \binom{\Delta-1}{3} L_{-3}^3 L_{-2}^{\Delta-4} \right) \Omega \\
& + (\xi_1 L_{-5} L_{-2}^{\Delta-2} + \xi_2 L_{-4} L_{-3} L_{-2}^{\Delta-3} + \xi_3 L_{-3}^3 L_{-2}^{\Delta-4}) \Omega . \tag{3.14}
\end{aligned}$$

Now comparing the coefficients of the vectors $L_{-5} L_{-2}^{\Delta-2} \Omega$, $L_{-4} L_{-3} L_{-2}^{\Delta-3} \Omega$ and $L_{-3}^3 L_{-2}^{\Delta-4} \Omega$ in (3.13) and (3.14) yields

$$\xi_1 = \frac{1}{2} \left(6B + C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} (\Delta-1) \right) , \tag{3.15a}$$

$$\xi_2 = \frac{1}{2} \left(-9B + 2B\Delta + C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} (\Delta^2 - 3\Delta + 2) \right) , \tag{3.15b}$$

$$\xi_3 = \frac{1}{24} \left(45B - 15B\Delta + C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} (2\Delta^3 - 12\Delta^2 + 22\Delta - 12) \right) . \tag{3.15c}$$

With this knowledge of both vectors $N_{-2\Delta}^{aa} \Omega$ and $N_{-2\Delta-1}^{aa} \Omega$ at length $\Delta - 1$, now one last piece of information can be utilized in order to find another way to compute B . Until now, only the quasi-primarity of the null-field has been used. But actually it is also primary, i.e. the relation $[L_m, N_n^{aa}] = ((2\Delta-1)m-n)N_{m+n}^{aa}$ holds for all $m, n \in \mathbb{Z}$. In particular, this is true for $m = 2$ and $n = -2\Delta - 1$, and thus

$$\begin{aligned}
0 &= (6\Delta-1)N_{-2\Delta+1}^{aa} \Omega = [L_2, N_{-2\Delta-1}^{aa}] \Omega = L_2 N_{-2\Delta-1}^{aa} \Omega \\
&\stackrel{\Delta-1}{=} -\frac{3}{2} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \frac{2\Delta-1}{4\Delta-3} (7(\Delta-1) + 6(\Delta-1)(\Delta-2)) L_{-3} L_{-2}^{\Delta-2} \Omega \\
&\quad + (7\xi_1 + 6\xi_2) L_{-3} L_{-2}^{\Delta-2} \Omega + [L_2, N^{(\Delta)}(W^a, W^a)_{-2\Delta-1}] \Omega \\
&\stackrel{\Delta-1}{=} -\frac{3}{2} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \frac{2\Delta-1}{4\Delta-3} (7(\Delta-1) + 6(\Delta-1)(\Delta-2)) L_{-3} L_{-2}^{\Delta-2} \Omega \\
&\quad + (7\xi_1 + 6\xi_2) L_{-3} L_{-2}^{\Delta-2} \Omega + [L_2, W_{-\Delta}^a W_{-\Delta-1}^a + W_{-\Delta-1}^a W_{-\Delta}^a] \Omega \\
&\stackrel{\Delta-1}{=} -\frac{3}{2} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \frac{2\Delta-1}{4\Delta-3} (7(\Delta-1) + 6(\Delta-1)(\Delta-2)) L_{-3} L_{-2}^{\Delta-2} \Omega \\
&\quad + (7\xi_1 + 6\xi_2) L_{-3} L_{-2}^{\Delta-2} \Omega \\
&\quad + ([L_2, W_{-\Delta}^a] W_{-\Delta-1}^a + [L_2, W_{-\Delta-1}^a] W_{-\Delta}^a) \Omega \\
&\stackrel{\Delta-1}{=} -\frac{3}{2} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \frac{2\Delta-1}{4\Delta-3} (7(\Delta-1) + 6(\Delta-1)(\Delta-2)) L_{-3} L_{-2}^{\Delta-2} \Omega
\end{aligned}$$

$$\begin{aligned}
& + (7\xi_1 + 6\xi_2) L_{-3} L_{-2}^{\Delta-2} \Omega + (2(\Delta - 1) + \Delta) [W_{-\Delta+2}^a, W_{-\Delta-1}^a] \Omega \\
& + (2(\Delta - 1) + \Delta + 1) [W_{-\Delta+1}^a, W_{-\Delta}^a] \Omega \\
\stackrel{\Delta-1}{=} & -\frac{3}{2} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} \frac{2\Delta - 1}{4\Delta - 3} (7(\Delta - 1) + 6(\Delta - 1)(\Delta - 2)) L_{-3} L_{-2}^{\Delta-2} \Omega \\
& + (7\xi_1 + 6\xi_2) L_{-3} L_{-2}^{\Delta-2} \Omega \\
& + (3\Delta - 2) C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} p_{\Delta, \Delta, 2\Delta-2}(2 - \Delta, -\Delta - 1)(\Delta - 1) L_{-3} L_{-2}^{\Delta-2} \Omega \\
& + (3\Delta - 1) C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} p_{\Delta, \Delta, 2\Delta-2}(1 - \Delta, -\Delta)(\Delta - 1) L_{-3} L_{-2}^{\Delta-2} \Omega, \quad (3.16)
\end{aligned}$$

where in this case the term $N^{(\Delta)}(W^a, W^a)_{-2\Delta-1} \Omega$ (recall equation (2.7)) *does* lead to a contribution at length $\Delta - 1$, in contrast to the situation in equation (3.14). Now using (3.12) and (3.15) in (3.16) yields the following alternate expression for the parameter B :

$$B = -\frac{12\Delta^2 - 18\Delta + 7}{4(4\Delta - 3)} C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})}.$$

This can only be in agreement with (3.11) for $C_{W^a, W^a}^{\mathcal{N}(T^{\Delta-1})} = 0$, which is not the case. Thus, the assumption $\alpha = 0$ leads to a contradiction and V_Δ is C_2 -cofinite.

We summarize our results in the following theorem.

Theorem. For all $p \in \mathbb{Z}_{\geq 2}$, the nonmeromorphic operator product expansion exists and is associative for the vertex operator algebra $\mathcal{W}(2, (2p-1)^{\times 3})$. Furthermore, all these vertex operator algebras are C_2 -cofinite.

References

- [A] T. Abe, *A \mathbb{Z}_2 -orbifold model of the symplectic fermionic vertex operator superalgebra*, [[math.QA/0503472](#)].
- [ABD] T. Abe, G. Buhl, and C. Dong, *Rationality, Regularity, and C_2 -cofiniteness*, *Trans. Amer. Math. Soc.* **356** (2004), 3391–3402, [[math.QA/0204021](#)].
- [BFKNRV] R. Blumenhagen, M. Flohr, A. Kliem, W. Nahm, A. Recknagel, and R. Varnhagen, *W-Algebras with two and three Generators*, *Nucl. Phys. B* **361** (1991), 255–289.
- [B] G. Buhl, *A spanning set for VOA modules*, *J. Alg.* **254** (2002), 125–151, [[math.QA/0111296](#)].
- [DLM] C. Dong, H. Li, and G. Mason, *Modular invariance of trace functions in orbifold theory*, *Commun. Math. Phys.* **214**, (2001), 1–56, [[q-alg/9703016](#)].

- [FHL] I. B. Frenkel, Y.-Z. Huang, and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Memoirs Amer. Math. Soc.* **104** (1989).
- [F1] M. Flohr, W-Algebren, Quasiprimäre Felder & Nicht-Minimale Modelle (1991), diploma thesis.
- [F2] M. Flohr, On modular invariant partition functions of conformal field theories with logarithmic operators, *Int. J. Mod. Phys. A* **11** (1996), 4147–4172, [[hep-th/9509166](#)].
- [F3] M. Flohr, Bits and pieces in logarithmic conformal field theory, *Int. J. Mod. Phys. A* **18** (2003), 4497–4592, [[hep-th/0111228](#)].
- [GK] M. R. Gaberdiel and H. Kausch, A rational logarithmic conformal field theory, *Phys. Lett. B* **386** (1996), 131–137, [[hep-th/9606050](#)].
- [GN] M. R. Gaberdiel and A. Neitzke, Rationality, quasirationality and finite W -algebras, *Commun. Math. Phys.* **238** (2003), 305–331, [[hep-th/0009235](#)].
- [G] M. R. Gaberdiel, An algebraic approach to logarithmic conformal field theory, *Int. J. Mod. Phys. A* **18** (2003), 4593–4638, [[hep-th/0111260](#)].
- [HL1] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, I, *Selecta Mathematica* **1** (1995), 699–756, [[hep-th/9309076](#)].
- [HL2] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, II, *Selecta Mathematica* **1** (1995), 757–786, [[hep-th/9309159](#)].
- [HL3] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, III, *J. Pure Appl. Algebra* **100** (1995), 141–172, [[q-alg/9505018](#)].
- [HLZ] Y.-Z. Huang, J. Lepowsky, and L. Zhang, A logarithmic generalization of tensor product theory for modules for a vertex operator algebra, [[math.QA/0311235](#)].
- [H1] Y.-Z. Huang, A theory of tensor products for module categories for a vertex operator algebra, VI, *J. Pure Appl. Algebra* **100** (1995), 173–216, [[q-alg/9505019](#)].
- [H2] Y.-Z. Huang, Differential equations and intertwining operators, *Comm. Contemp. Math.* **7** (2005), 375–400, [[math.QA/0206206](#)].

- [Kad] L. P. Kadanoff, Operator algebra and the determination of critical indices, *Phys. Rev. Lett.* **23** (1969), 1430–1433.
- [Kau] H. G. Kausch, Extended conformal algebras generated by a multiplet of primary fields, *Phys. Lett. B* **259** (1991), 448–455.
- [L] H. Li, Some finiteness properties of regular vertex operator algebras, *J. Alg.* **212** (1999), 495–514, [[math.QA/9807077](#)].
- [LL] J. Lepowsky and H. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Mathematics, Birkhäuser, 2004.
- [M] M. Miyamoto, A theory of tensor products for vertex operator algebra satisfying C_2 -cofiniteness, [[math.QA/0309350](#)].
- [N] W. Nahm, *Conformal Quantum Field Theory in Two Dimensions*, unpublished.
- [R] F. Rohsiepe, On Reducible but Indecomposable Representations of the Virasoro Algebra, *BONN-TH-96-17* (1996), [[hep-th/9611160](#)].
- [W] K. G. Wilson, Non-Lagrangian models of current algebra, *Phys. Rev.* **179** (1969), 1499–1512.
- [Z] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996), 237–302.